



California Greenhouse Gas Emission Measurement Project (CALGEM)

M.L. Fischer, C. Zhao, W.J. Riley,
Lawrence Berkeley National Laboratory

A. Andrews, A.I. Hirsch, C. Sweeny, P. Tans
NOAA ESRL Global Monitoring Division

Acknowledgements: P. Bousquet, J.C. Lin, C. Potter,
W. Salas, T. Szegvary. This work was supported by the
California Energy Commissions PIER program.


Contact: MLFischer@LBL.gov

Outline

- CH₄ and non-CO₂ GHGs
 - Importance, uncertainty
 - Need for multiple emissions constraints
- Design of a regional measurement network
 - Emission inventory maps
 - Tower sites and instrumentation
 - Estimation of atmospheric signals
- Summary
- Further work


Overview of non-CO₂ GHGs

- Total non-CO₂ GHG forcing is ~ equivalent to CO₂ forcing globally
 - CH₄N₂O, High GWP
 - Tropospheric ozone
 - GHG feedbacks to climate change are likely (How big?)
- California non-CO₂ emissions:
 - equivalent to ~ 15% of CO₂
 - Large uncertainties because sources not readily metered
 - Currently, data are lacking to accurately scale up to state level
 - Alternately, look from top down using atmosphere for integration



Emissions (MMTCO₂eq)

CA non-CO₂ Emissions, 1999




Constraining GHG Emissions with Atmospheric Data-Model Synthesis

- Essential ingredients:
 - A priori inventory estimates for spatiotemporal distribution of GHG and other tracers
 - Model for atmospheric transport and surface influence “footprints”
 - Continuous long term measurements of GHG and source attribution species (e.g., isotopes, atmos. tracers)
 - Quantitative GHG boundary conditions for regional problem
 - Bayesian statistical framework to evaluate improvement in emission estimates

A priori CH₄ Emission Inventories


- Landfill
 - Landfill specific loading with substrate dependent residence time (EPA)
- Animal Ag.
 - USDA county level stocking
 - Dairy/meat emission factor
- Natural gas dist./use
 - County level facility/usage statistics (ARB)
- Wetlands
 - NASA-CASA (Potter, 2006)
- Crop Agriculture (July)
 - County level DNDC (Salas et al., 2006)



Atmospheric Mixing and Source Attribution

- ^{222}Rn can constrain atmospheric mixing (Hirsch, 2007)
 - Nearly ubiquitous soil emission
 - Atmospheric concentrations reflect recent emissions because of short half life (3.8 day)
 - Soil moisture is a factor
- Isotopic signatures
 - $^{13}\text{CO}_2$ separate nat. gas vs. gasoline (Pataki, 2006)
 - ^{14}C not present in fossil fuel
 - $^{13}\text{CH}_4$, CDH_3 separate landfills and nat. gas
- CO, VOCs (specific to animals, crops, petroleum, etc.)

Soil ^{222}Rn Emissions




(Szegvary, 2006)

Initial Measurement Sites




**Oceanic Site:
Sutro Tower,
San Francisco
(116, 230 m agl)**



**Inland Site:
KCRA Tower,
Walnut Grove
(30,100, 480 m agl)**

Measurement Instrumentation


- 12 Flask System
 - Twice daily samples
 - $\text{CO}_2, \text{CH}_4, \text{N}_2\text{O}, \text{CO}$
 - SF_6 , halo carbons
 - $^{13}\text{CO}_2, ^{13}\text{CH}_4, \text{CDH}$
- CH_4/CO_2 analyzer- 3min
- CO_2/CO rack – 3 min
- ^{222}Rn monitor – 30 min
- $^{14}\text{CO}_2$ (w/ LLNL soon)
- Collaborations sought



Predicting which Land Surface Affects the Measurements: The Footprint

- Stochastic time inverted Lagrangian particle trajectory (STILT) model predicts source regions contributing to measured signals
- Requires mesoscale model (e.g., NOAA-EDAS, MM5/WRF) for winds, turbulence, convection

Area affecting concentrations at Sutro Tower (232 m), July 2005





Predicted CH₄ Signals

- Signals at Sutro dominated by background oceanic air
 - ²²²Rn peaks show periods with influence of land surface
- Signals at KCRA dominated by emissions from central valley
 - Strong diurnal cycle from changing boundary layer height
 - Synoptic variations also apparent

Sutro tower at 232 m


KCRA at 100 m



CH₄ vs. ²²²Rn Signals

KCRA at 100 m

- CH₄ signals correlated with ²²²Rn
 - ²²²Rn acts as a tracer of land surface influence
 - Measurements combined with quantitative footprints provides constraint on GHG emissions




Summary

- California's non-CO₂ GHG emissions significant and uncertain
- Long term atmospheric measurements will provide an independent and complementary method to constrain estimates of regional emissions
- Initial prediction of CH₄ signals suggest measurements can be achieved with planned instrumentation

Further Work


- Initiate measurements at Sutro and Walnut Grove Towers
- Develop and test high resolution meteorology for tower sites using MM5 and Weather Research Forecast (WRF) model
- Incorporate boundary conditions, isotopic tracers, and species for source attribution
- Initiate model-data-synthesis of regional GHG emissions and their uncertainties

A wide-angle photograph of a sunset. The sun is a bright, glowing orb on the horizon, partially obscured by a range of dark mountains. The sky is filled with layers of clouds, which are illuminated from below, creating a vibrant palette of orange, yellow, and red. The lower portion of the image shows a dark, silhouetted foreground of trees and a cityscape. In the distance, across a body of water, the Golden Gate Bridge is visible on the left side. The overall mood is peaceful and majestic.

Thank you

Example Transport Simulation: Atmospheric Signals of Fossil Fuel CO₂


- Forward model simulation
 - Fossil CO₂ emissions scaled from EPA-NO_x
 - MM5-Tracer
 - 10km resolution
 - Smaller nested grids possible over sub-regions of interest
 - Plumes from urban areas transport into surrounding regions (Riley et al., 2007)



Global Assimilated CH_4 Background

- Oceanic values indicate background air arriving at California
- Global assimilation results valuable for subtraction from tower “signals”
- Latitudinal gradients and long distance transport likely important
- Coastal measurements also needed to capture unresolved variations in background CH_4

Monthly mean CH_4 for 2003



Bousquet et al., (2004)